



## Where is the new physics?

Muon anomalous magnetic moment (g-2) sensitive to quantum-mechanical effects of new heavy or weakly-coupled particles (e.g SUSY, extra dimensions, dark photons, ...)




- ♦ Measured to very high precision (0.54 ppm!)
  - → provides some of strongest constraints on extensions of Standard Model
- ♦ BNL measurement disagrees with Standard-Model theory expectations by >3σ




## Motivation: the Muon g-2 Experiment




- ◆ Fermilab Muon g-2 Experiment aims to reduce BNL measurement error by four
  - Started running in June, and expects first BNL-level results in 2018!
- → Must reduce theory error to commensurate level to identify definitively whether any deviation observed is due to new physics




### Muon g-2 in the Standard Model






#### Hadronic contributions to muon g-2

# **Hadronic Vacuum Polarization (HVP)**



- ◆ Obtained from experimental measurements for e<sup>+</sup>e<sup>-</sup>→hadrons plus dispersion relations
- ◆ challenging analysis combining more
  than 20 exclusive channels with several
  final-state hadrons [Davier et al.,
  Keshavarzi et al., Jegerlehner]


#### **Hadronic Light-by-Light (HLbL)**



- ◆ Estimated from QCD models such as large N<sub>c</sub>, chiral perturbation theory, vector meson dominance, ...
- ◆ Subjective and not systematically improvable [Glasgow consensus, Prades, de Rafael, Vainshtein, 0901.0306]


#### Hadronic contributions to muon g-2

# **Hadronic Vacuum Polarization (HVP)**



Obtained from experimental

#### **Hadronic Light-by-Light (HLbL)**




- ◆ Estimated from OCD models such as
- ◆ To match anticipated experimental precision, must reduce HVP uncertainty to below ~0.2% and HLbL uncertainty to below ~10%
- → Both contributions are calculable (in principle) from ab-initio Quantum chromodynamics with numerical lattice QCD!

Keshavarzi et al., Jegerlehner

0901.0306]

#### Numerical lattice Quantum Chromo Dynamics

- Systematic method for calculating hadronic parameters from QCD first principles
- Define QCD on (Euclidean) spacetime lattice and solve path integral numerically
  - **♦** Recover QCD when lattice spacing  $a \rightarrow 0$  and box size L→∞




- Simulate using Monte-Carlo methods and importance sampling
  - ❖ Sample from all possible field configurations using a distribution given by exp(-S<sub>QCD</sub>)
- Run codes upon supercomputers and dedicated clusters




## 21st-century lattice QCD

- Modern lattice-QCD methods + latest generations of tera/petascale supercomputers have enabled calculations of many simple quantities with controlled uncertainties and complete error budgets
- Standard simulations include dynamical
   u, d, & s quarks in the vacuum
  - Typically isospin symmetric (mu=md)
  - Many now include dynamical charm
- ◆ Control systematic errors using QCD gauge-field ensembles with different parameters (quark masses, lattice spacings, spatial volumes, ...)
  - State-of-the-art calculations employ pions with physical mass ~135 MeV!



#### Lattice-QCD validation

- ◆ Lattice-QCD results agree with experimental measurements for wide variety of hadron properties including light- and heavy-hadron spectrum & proton-neutron mass difference
- Lattice-QCD predictions
   of B<sub>c</sub> mass, D<sub>(s)</sub> decay
   constants, & D→KIv form
   factor appeared before
   measurements with
   comparable precision
- ◆ Independent lattice-QCD calculations using different actions & methods provide confirmation for matrix elements inaccessible by experiment
- ◆ Demonstrate that calculations are reliable with controlled errors!





#### **Participants**

We are using *ab-initio* lattice-QCD to target the hadronic vacuum-polarization contribution, which is the largest source of theory error



- Aida El Khadra (Illinois)
- Andreas Kronfeld (Fermilab)
- Ethan Neil (Colorado)
- Ruth Van de Water (Fermilab)

#### **MILC** Collaboration

- Carleton DeTar (Utah)
- Steve Gottlieb (Indiana)
- Jack Laiho (Syracuse)
- Yuzhi Liu (Indiana)



- Bipasha Chakraborty (JLAB)
- Daniel Hatton (Glasgow)
- Christine Davies (Glasgow)
- Jonna Koponen (INFN, Rome)
- Peter Lepage (Cornell)
- Andrew Lytle (Glasgow)
- Craig McNeile (Plymouth)

Doug Toussaint (Arizona)

Alejandro Vaquero (Utah)

#### HVP contribution from lattice QCD

[Blum, PRL91 (2003) 052001]

◆ Lowest-order hadronic-vacuum polarization contribution obtained from 1-loop QED

integral:

- ❖ Kinematic function f(Q²) accounts for interaction between muon and external photon
- \* Vacuum polarization function  $\Pi(Q^2)$  captures nonperturbative QCD effects
- ❖  $\Pi(Q^2)$  related to inclusive e<sup>+</sup>e<sup>-</sup>→hadrons cross section via dispersion relation
- $\curvearrowright$  Can also compute Euclidean space  $\Pi(Q^2)$  directly in lattice QCD from simple two-point correlation function of electromagnetic quark current
  - $\clubsuit$  In principle can yield more precise  $\prod(Q^2)$  unlimited by experimental uncertainties

### Π(Q²) from moments + Padé approximants


[HPQCD, PRD89, no. 11, 114501 (2014)]

- ❖ New method introduced by HPQCD enables precise determination of vacuum polarization function  $\Pi(Q^2)$  from zero-momentum vector-current correlation functions:
- (1) Calculate derivatives of vacuum polarization function  $\Pi(Q^2)$  at  $Q^2=0$  from time moments of vector-current correlators

$$\hat{\Pi}(Q^2) = \sum_{j=1}^{\infty} Q^{2j} \Pi_j$$

$$\Pi_j = \frac{(-1)^{j+1}}{(2j+2)!} \sum_t t^{2j+2} \langle j_{\text{EM}}^{\mu}(t) j_{\text{EM}}^{\mu}(0) \rangle$$

(2) Use Taylor coefficients to construct [n,n] and [n,n-1] Padé approximants for  $\Pi(Q^2)$ 



◆ Demonstrate method with s-,c-, and b-quark connected contributions [PRD 89, no. 11, 114501 (2014); PRD 91, no. 7, 074514 (2015)]

#### 2016 result for u/d-connected contribution


- ◆ Employ large set of four-flavor QCD gauge-field ensembles with dynamical u,d,s, & c quarks [MILC Collaboration, PRD82 (2010) 074501]
  - Three lattice spacings, multiple spatial volumes, physical light-quark masses to control systematic uncertainties
  - Largest errors from omission of isospin breaking and electromagnetism
  - Errors from the nonzero lattice spacing and finite spatial volume are next

$a_{\mu}^{\mathrm{HVP,LO}}(u/d)$
$\frac{a_{\mu}}{1.0\%}$
1.0%
0.7%
0.4%
0.4%
0.4%
0.3%
0.4%
0.2%
0.2%
< 0.05%
1.8%

[HPQCD + RV, 1601.03071; to be published in PRD]

#### Total LO HVP contribution: 2016


#### **Contributions by quark flavor**



 $10^{10} a_{\mu}^{\text{HVP,LO}} = 667(11)_{u,d}(1)_{s,c,b}(9)_{\text{disc.}}$ 


[HPQCD + RV, 1601.03071; to be published in PRD]

First complete lattice-QCD calculation of leading-order  $a_{\mu}^{HVP}$  to reach precision needed to observe significant deviation from experiment



## Fermilab/HPQCD/MILC work in progress


- Fermilab Lattice, HPQCD, & MILC Collaborations joined efforts for followup calculation to reduce error and meet target experimental precision
- ♦ Addressing leading sources of error from:
  - (1) Omission of quarks' EM charges → developing code to generate new ensemble with dynamical photons
  - (2) u/d-quark mass difference →
    generated new 1+1+1+1 ensemble
    (see next slide)
  - (3) Quark-disconnected contributions → implemented modern variance-reduction algorithms in MILC code & testing parameters
  - (4) Finite spatial volume & nonzero lattice spacing → analyzing ensembles with finer lattice spacings




Aim to reach ~1% uncertainty before first result from Muon g-2 Experiment next year

## New! Isospin-breaking correction

- First result from joint effort is direct calculation of m<sub>u</sub>≠m<sub>d</sub> correction using new ensemble with non-degenerate up and down sea quarks
- ♦ Fix average  $(m_u+m_d)/2 =$  27.35(11) and ratio  $m_u/m_d$  = 0.4582 to reproduce physical  $M_{π0}$  and electromagnetic splitting  $(M_{π+} M_{π0})$  [MILC, PRD90 (2014) no.7, 074509;1606.01228]
- First lattice-QCD calculation of the strong-isospin-breaking contribution to a<sub>µ</sub> HVP,LO at the physical pion mass





## Progress on hadronic light-by-light


- ◆ Most recent lattice calculations combine nonperturbative QCD gauge-field configurations with perturbative QED for the photons to directly obtain desired O(a<sub>EM</sub><sup>3</sup>) contribution
- ♦ New method from RBC Collaboration employs stochastically-generated photon propagators to control statistical noise [Blum et al., PRD93, 014503 (2016); PRL118 (2017) no.2, 022005]
  - ♦ Obtain ≤10% statistical errors at the physical pion mass in ballpark of Glasgow consensus value  $a_{\mu}^{HLbL,GC} \times 10^{10} = 10.5(2.6)$


$$a_{\mu}^{\rm HLbL} \times 10^{10} = \begin{cases} 11.60(0.96)_{\rm stat.} & \rm connected \\ -6.25(0.80)_{\rm stat.} & \rm disconnected \end{cases}$$

- Full study of systematic errors (especially finite-volume effects) needed, but initial results encouraging!
- ◆ Other complementary approaches being studied by Mainz group:
  - Combination of lattice QCD for nonperturbative 4-point function + continuum, infinite-volume perturbation theory for QED kernel [Asmussen et al., 1609.08454]
  - ♦ Calculation of dominant pion-pole contribution using lattice QCD for  $\pi^0 \rightarrow \gamma^* \gamma^*$  form factor + dispersion relation [Géradin et al., PRD94 (2016) no.7, 074507]

#### Quark-disconnected contributions to HVP


- ◆ Contributions in which valence quarks are connected by a pair of gluons only few percent of total HVP contribution due to SU(3)<sub>flavor</sub> and Zweig suppression
- ◆ Computationally challenging due to large statistical noise, but now possible to obtain statistically-significant signal using modern variance-reduction strategies such as allmode averaging [Blum et al, PRD 88 (2013) 094503]
- ◆ First calculation of quark-disconnected contribution with physical-mass pions last year by RBC/UKQCD [PRL116, 232002 (2016)]
- ♦ Preliminary BMW Collaboration result with higher statistics has uncertainty  $\delta a_{\mu}^{HVP,disc.}$  ~ 1.5×10<sup>10</sup> → contributes only 0.2% to error on total  $a_{\mu}^{HVP,LO}$






### Hybrid lattice+experiment approach for HVP

- Spectral representation enables comparison of vacuum polarization function from lattice QCD with determination from experimental e<sup>+</sup>e<sup>-</sup>→hadrons ("R-ratio") data [Bernecker & Meyer, EPJA47 (2011) 148, arXiv:1107.4388]
  - ♦ Most precise in complementary Q² regions → combination yields smaller error than lattice QCD or experiment alone
  - $\Rightarrow$  May provide more immediate path to target experimental precision until first-principles lattice-QCD-calculations reach this goal






[C. Lehner, Lattice 2017]



#### Summary & outlook

- ◆ Lattice-QCD community is working hard to compute hadronic contributions to muon g-2 to the precision needed by the Muon g-2 (and planned J-PARC) Experiment
- **→** Methods in place for hadronic vacuum polarization contribution
  - Several independent calculations ongoing using different approaches
  - First results reported for previously omitted effects of isospin breaking & electromagnetism and contributions from quark-disconnected diagrams
  - Improvements to most precise calculation by HPQCD+RV in progress by Fermilab, HPQCD, and MILC Collaborations
  - \* Hope to reach sub-per cent precision from lattice QCD in 1 or 2 years!
- ◆ Methods being developed and tested for hadronic light-by-light contribution
  - Study of systematic errors in promising RBC-Collaboration approach in progress
  - Still room for new theoretical developments to make substantial impact



### Muon g-2 Theory Initiative

- ◆ GOAL: provide theoretical predictions for hadronic contributions to g-2 with reliable & precise uncertainties to maximize impact of Fermilab & J-PARC experiments
- ◆ PLAN: organize series of workshops bringing together theorists from lattice & non-lattice communities to:
  - Discuss, assess, & compare efforts and approaches
  - Map out strategies for improving theoretical calculations
  - Write reports summarizing status of relevant theoretical work
- TIMELINE: finalize first report in September 2018 before Muon g-2 Experiment announces first BNL-level measurement



★ Kickoff workshop hosted by Fermilab in June with 66 registered participants, 40 talks in 15 discussion sessions over 4 days




### HVP from e⁺e⁻→hadrons

♦ Standard-Model value for  $a_{\mu}^{HVP}$  obtained from experimental measurement of  $\sigma_{total}(e^+e^- \rightarrow hadrons)$  via optical theorem:

$$a_{\mu}^{\text{HVP}} = \left(\frac{\alpha m_{\mu}}{3\pi}\right)^2 \int_{m_{\pi^0}^2}^{\infty} ds \frac{R(s)K(s)}{s^2} \qquad R \equiv \frac{\sigma_{\text{total}}(e^+e^- \to \text{hadrons})}{\sigma(e^+e^- \to \mu^+\mu^-)}$$


- (Away from quark thresholds, use four-loop pQCD)
- ◆ Includes >20 multi-particle channels with up to six final-state hadrons
- Multi-hadron channels represent a small absolute contribution to a<sub>µ</sub><sup>HVP</sup>, but contribute a significant fraction of the total uncertainty




[Jegerlehner and Nyffeler, Phys.Rept. 477 (2009) 1-110]

## HLbL from QCD models

- Hadronic light-by-light contribution cannot be expressed in terms of experimental quantities and must be obtained from theory
   [cf. Jegerlehner and Nyffeler, Phys.Rept. 477 (2009) 1-110 and Refs. therein]
  - ❖ All recent calculations compatible with constraints from large-N<sub>c</sub> and chiral limits
  - \* All normalize dominant  $\pi^0$ -exchange contribution to measured  $\pi^0 \rightarrow \gamma \gamma$  decay width
  - Differ for form factor shape due to different QCD-model assumptions such as vector-meson dominance, chiral perturbation theory, and the large N<sub>c</sub> limit





# The Glasgow consensus for HLbL

- ♦ Quoted error on for a<sub>μ</sub><sup>HLbL</sup> from Prades, de Rafael, Vainshtein, 0901.0306 based range on model estimates
  - $\pi^0$ -exchange contribution estimated to be ~10 times larger than others
  - ❖ Largest contribution to uncertainty (±1.9×10<sup>-10</sup>) attributed to charged pion and kaon loop contributions

Table 1: Contribution to  $a^{\text{HLbL}}$  from  $\pi^0$ ,  $\eta$  and  $\eta'$  exchanges

Result	Reference
$(8.5 \pm 1.3) \times 10^{-10}$	[7, 8]
$(8.3 \pm 0.6) \times 10^{-10}$	[4, 5, 6]
$(8.3 \pm 1.2) \times 10^{-10}$	[1]
$(11.4\pm1.0)\times10^{-10}$	[9]



$$a^{\rm HLbL}(\pi\,,\eta\,,\eta') = (11.4 \pm 1.3) \times 10^{-10}$$

 Subjective error estimate does not cover spread of model values, and and is not systematically improvable